

Analysis of Disease by Image Segmentation Using Retinal Blood Vessels Detection

Kodi Suma

Assistant Professor, ECE, Dadi Institute of Engineering & Technology, India.

B.T. Archana

Assistant Professor, ECE, Dadi Institute of Engineering & Technology, India.

Abstract – A retinal vessel occlusion is a blockage in the blood vessel of your eye that can result in sight loss. Eye ailments and sight loss are one of the significant health issues among the senior and old individuals. Retina is the most important internal components in eye. Retinal images play vital role in several applications such as disease diagnosis and human recognition. They also play a major role in early detection of diabetics by comparing the states of the retinal blood vessels. Retinal Image Analysis is a key element in detecting retinopathies in patients. Diabetic Retinopathy is one of the most common diabetic eye conditions which cause blindness induced by alterations in the blood vessels of the retina. In this work we detect the blood vessels effectively for the diagnosis of disease such as diabetes, glaucoma, and hemorrhage.

Index Terms – Retinal vessel occlusion, Retinopathy, Blood vessels, Retinal image, Segmentation, Diagnosis.

1. INTRODUCTION

Human eye is an important organ that reacts to light and has several purposes. The eye has a number of components which include but are not limited to the cornea, iris, pupil, lens, retina, and macula, optic nerve, choroid and vitreous.

The eye is not shaped like a perfect sphere; rather it is a fused two-piece unit. The smaller frontal unit, transparent and more curved, called the cornea is linked to the larger white unit called the sclera. The eye is made up of three coats:

- **Outermost layer-** The outermost layer, known as the fibrous tunic, is composed of the cornea and sclera.
- **Middle layer-** The middle layer, known as the vascular tunic or uvea, consists of the choroid, ciliary body, and iris.
- **Innermost layer -** The innermost is the retina, which gets its circulation from the vessels of the choroid as well as the retinal vessels, which can be seen in an ophthalmoscope.

The retina is internal part of the eye. In the center of retina there is the optic disk, a circular to oval shape. From the center of optical nerve radiates the major blood vessels of the retina. The blood vessels network is an important anatomical

structure in human retina, which is used to recognize different types of disease. There are two types of retinal blood vessels, arteries and veins. However, manual detection of blood vessels is not simple because the vessels in retina image are complex and have low contrast. For retinal anatomy ophthalmologist uses an ophthalmoscope. The retinal fundus image is widely used in the diagnosis and treatment of various types of diseases such as diabetic retinopathy and glaucoma. There are different types of eye diseases, such as Cataract, Iridocyclitis, Corneal Haze, Glaucoma and Diabetic retinopathy.

Here we see the three types of disease description

- **Diabetes:** Diabetes is a long-term condition that causes high blood sugar levels. Diabetes is a serious complex condition which can affect the entire body? When someone has diabetes, their body can't maintain healthy levels of glucose in the blood. Glucose is a form of sugar which is the main source of energy for our bodies.
- **Glaucoma:** Glaucoma is a group of eye diseases which in most cases produce increased pressure within the eye- if left untreated the patient may lose vision and even become blind. This elevated pressure is caused by a backup of fluid in the eye.
- **Hemorrhage:** Is a disorder of the eye in which bleeding occurs in the light sensitive tissue on the back wall of the eye. A retinal hemorrhage can be caused by hypertension, retinal vein occlusion (a blockage of a retinal vein), or diabetes mellitus (which causes small fragile blood vessels to form, which are easily damaged).

Retinal blood vessels share similar size, structure, and function with blood vessels in the brain and can provide a way of examining brain health in living humans. The retina is a light-sensitive layer at the back of the eye that covers about 65 percent of its interior surface. Photosensitive cells called rods and cones in the retina convert incident light energy into signals that are carried to the brain by the optic nerve. A

retinal hemorrhage can be caused by hypertension, retinal vein occlusion (a blockage of a retinal vein), or diabetes mellitus (which causes small fragile blood vessels to form, which are easily damaged). However, some retinal hemorrhages can cause severe impairment of vision.

Checking the obtained changes in retinal images in an especial period can help the physician to diagnose the disease. Applications of retinal images are diagnosing the progress of some cardiovascular diseases, diagnosing the region with no blood vessels (Macula). Retinal image analysis is a complicated task particularly because of the variability of the images in terms of the color, the morphology of the retinal anatomical pathological structure and the existence of particular features in different patients, which may lead to an erroneous interpretation.

• Motivation

Detection of retinal blood vessels for disease diagnosis has provided more information about retinal blood vessels and disease. Compared with the other more traditional technology, detection of retinal blood vessels for disease diagnosis has the benefits of high anti-counterfeiting strength, small imaging devices, low cost, easy collection of images with contactless operation universality and liveliness. Furthermore, since the blood vessels are located internally within the living body, the disease identification system is less affected by the outer skin surroundings (skin disease, humidity, dirtiness, etc.). Hence retinal blood vessels for disease identification are considered as one of the most promising solution for disease diagnosis in the future.

Diabetic retinopathy affects the human eye and causes the blindness. For the efficient diagnosis of retinopathy accurate measurement (true vessel structure) of vessel diameter is required for estimating the types of vessels [1]. True approximation of total number of microaneurysms is required for estimating the stages of diabetic retinopathy. One of the most common disease taking place is diabetic retinopathy, which cause effect on eye, it top cause of loss of sight in the developed countries .One of the most important attribute of this illness is existence of microaneurysms (MAs) on retina. The presence of microaneurysms in eye is one of the early sign of diabetic retinopathy, which is one of leading cause of loss of vision [2]. The size of microaneurysms range from 10–100 microns which means these are less than 1/12 of diameter of the optic disc [3].

2. LITERATURE REVIEW

[4]Archna Sharma and Hempriya have proposed a method for Detection of Blood Vessels and Diseases in Human Retinal Images. In this the detection of blood vessels is important task in diagnosis the diseases of eye. The present study is aimed at developing an automated system for the extraction of normal

and abnormal features in retinal images. The blood vessel network is an important anatomical structure in human retina. Several diseases such as Diabetic retinopathy, glaucoma, hemorrhages, the performance of automatic detection methods may be improved if regions containing vessels can be excluded from the analysis.

[5] K.Jeyasri has proposed method on the detection of the retinal images and disease diagnosis. In this paper Retinal images play vital role in several applications such as disease diagnosis and human recognition. In this work a new algorithm to detect the blood vessels effectively has been proposed. Initially enhancement of the image is carried out using curvelet transform and modification of the curvelet coefficients.

[6] D. J. Cornforth and H. J. Jelinek have proposed Development of retinal blood vessel segmentation methodology using wavelet transforms for assessment of diabetic retinopathy. Automated image processing has the potential to assist in the early detection of diabetes, by detecting changes in blood vessel diameter and patterns in the retina. This paper describes the development of segmentation methodology in the processing of retinal blood vessel images obtained using non-mydratic colour photography. The methods used include wavelet analysis; they show highly accurate identification of blood vessels for the purpose of studying changes in the vessel network that can be utilized for detecting blood vessel diameter changes associated with the pathophysiology of diabetes.

[7]VijayaR.Patil proposed Detection of Optic Disc in Retina Using Digital Image Processing. The retinal fundus image is widely used in the diagnosis and treatment of various eye diseases such as diabetic retinopathy and glaucoma. He propose a method to automatically detect the optic disc in fundus images of the retina. The method includes edge detection using the canny operators and detection of circles using the Hough transform method.

[8]Mohammadreza Yadollahi published paper on “Image segmentation for object detection” in this paper there is brief introduction of segmentation method. It is used in many scientific fields including medical imaging, objectand face recognition, engineering and technology.

[9]Reyhaneh Sadeghzadeh and Michael Berks have proposed Detection of Retinal Blood Vessels Using Complex Wavelet Transforms and Random Forest Classification. In this paper a new method for detecting vessels in retinograms. The Dual-tree Complex Wavelet Transform (DT-CWT) is used to provide a rich, multi-scale description of local structure, and a random forest classifier is used to classify pixels as vessel/non-vessel on the basis of their DT-CWT coefficients. The method is tested on retinograms obtained from a publicly

available database and our results are compared with previously reported results for the same database.

3. PROPOSED METHODOLOGY

- Image Acquisition

Vessels patterns, invisible to the naked eye, can be viewed through an image sensor sensitive to infrared light. Infrared light passes through the tissues of the human body and is blocked by pigments such as hemoglobin or melanin. As hemoglobin exists densely in blood vessels, infrared light shining through causes the vessels to appear as dark shadow lines in the captured image. In image acquisition module there are two cameras used one is CCD camera and the other is web camera.

Following figures shows the captured retinal blood vessels images.




Figure 1 Sample Retinal Blood Vessels Image




Figure 2 Sample Retinal Blood Vessels Image With Diabetes

- Image Segmentation

Image segmentation is the authorized process of dividing an image into multiple parts. This is typically used to identify objects or other relevant information in digital images. Image segmentation is the process of partitioning a digital image into multiple segments (sets of pixels, also known as super pixels). The goal of segmentation is to simplify or change the representation of an image into something that is more meaningful and easier to analyze.




Figure 3 Diseased Retina Image with Hemorrhage

Image segmentation is typically used to locate objects and boundaries (lines, curves, etc.) in images. More precisely, image segmentation is the process of assigning a label to every pixel in an image such that pixels with the same label share certain visual characteristics. The result of image segmentation is a set of segments that collectively cover the entire image or a set of contours extracted from the image. Each of the Image segmentation is typically used to locate objects and boundaries (lines, curves, etc.) in images.

More precisely, image segmentation is the process of assigning a label to every pixel in an image such that pixels with the same label share certain visual characteristics. The result of image segmentation is a set of segments that collectively cover the entire image or a set of contours extracted from the image. Each of the pixels in a region is similar with respect to some characteristic or computed property, such as color, intensity, or texture. The original image is captured with the black unwanted background. Including the background reduced the accuracy of the original image, because the position of retinal blood vessels usually varies across different retinal blood vessels images, it is necessary to image segmentation in region of interest (ROI) before feature extraction and matching with database.

- Image enhancement

The segmented retinal blood vessels image is then enhanced to improve its contrast as shown in Figure. The image is resized to 1/4 of the original size, and enlarged back to its original size.

Bicubic interpolation is used in this resizing procedure. Finally, histogram equalization is used for enhancing the gray level contrast of the image.

- Feature extraction

Feature extraction is most important step in retinal blood vessels recognition algorithm. It is a special form of dimensionality reduction. It is a transformation of input data into the set of features. In the feature extraction process the canny edge detection method is used for feature extraction process. Extraction of features such as edges and curves from an image is useful for final authentication. Edges are important features in an image, they represents significant local intensity changes as shown in the following fig.




Figure 4 Filter Image Of Blood Vessels

- Matching With Database

The retinal blood vessels matching module is use in MATLAB; the MATLAB is used to execute the retinal blood vessels recognition algorithm. In the working of real time proposed retinal blood vessels recognition algorithm contains two stages. Enrollment stage and verification stage.

4. RESULTS AND DISCUSSIONS

Following is the result that we get after applying proposed methodology to different images, and get the disease name. First we see the segmentation and histogram of the database images.

In this section all the results and the discussions should be made.




Figure 5 Segmentation of Image

After that we set an input image for the recognition of the disease and extract the feature as follows, then it matches with the database images and find the disease.




Figure 6 Input Image For Recognition Of Disease

The input is matches with the databases. Then the disease is diagnosis.




Figure 7 Detection of Glaucoma Disease

5. CONCLUSION

This paper provides analysis of disease using retinal blood vessels detection. Where we find out the disease such as diabetes, glaucoma, and hemorrhage on the basis of their segmentation. We use segmentation technique for retinal blood vessels detection for database images and input image then we find out the disease.

REFERENCES

- [1] Rahul Chauhan; Anita Uniyal; V. P Dubey Detection of retinal blood vessels and reduction of false microaneurysms for diagnosis of diabetic retinopathy 2016 International Conference on Emerging Trends in Communication Technologies (ETCT)
- [2] Ganesh Naga Sai Prasad V Habibulla Khan E. Gopinathan "Reduction of False Microaneurysms in Retinal Fundus Images using Fuzzy C-Means Clustering in terms NLM Anisotropic Filter" international journal of computer applications(0975-8887) vol. 113 no.19 march 2015
- [3] Meera walvekar Geeta Salunke "Detection of diabetic retinopathy with feature extraction using image processing" international journal of emerging technology and advanced engineering vol. 5 no. 1 January 2015.
- [4] Cemilkirbas and Francis Quek, "A REVIEW OF VESSEL EXTRACTION TECHNIQUES AND ALGORITHMS", ACM computing surveys, Vol. 36, No. 2, June 2004.
- [5] Seyed Mohsen Zabihi, MortezaDelgir, and Hamid Reza Pourreza, "Retinal Vessel Segmentation Using Color Image Morphology and Local Binary Patterns", IEEE 2010.
- [6] AshaGowdaKaregowda,AsfiyaNasiha,M.A.Jayaram, A.S. Manjunath, "Exudates Detection in Retinal Images using Back propagation Neural Network", vol. 25- No. 3, july 2011.
- [7] J. Benadict Raja, C.G. Ravichandran, "Blood Vessel Segmentation For High Resolution Retinal Images",IJCSI, Vol.8, Issue 6, No 2, November 2011
- [8] Shilpa Joshi, Dr. P.T. Karule, "Retinal Blood Vessel Segmentation", IJEIT, Vol.1, Issue 3, March 2012.
- [9] JaspreetKaur, Dr. H.P.Sinha, "An Efficient Blood Vessel Detection Algorithm for Retinal Images using Local Entropy Thresholding", (IJERT), ISSN: 2278-0181, Vol. 1 Issue 4, June 2012.
- [10] Seyed Mohsen Zabihi, Hamid Reza Pourreza, ToukaBanaee, "Vessel Extraction of Conjunctival Images Using LBPs and ANFIS", International Scholarly Research Network, ISRN Machine Vision, vol. 2012.

Authors



K.Suma born in 1991 in Gowripalli, AP. She pursued her B.Tech in GIET, Rajahmundry in 2012 and completed her Post Graduation in GIET, Rajahmundry in 2016. She has 5 years of teaching experience. Currently she is working as an assistant professor in DIET and her areas of interest are communications and image processing.



B.T.Archana was born in 1985 in Kerala. She pursued her B.Tech in Marian Engineering College in 2006 and completed her Post Graduation in 2014 in SCT college of Engineering in Kerala. She has 3 years of teaching experience. Currently she is working as an Assistant Professor in Dadi Institute of Engineering & Technology. Her areas of interest are Image Processing and signal processing.